

# Causal Inference on Multivariate and Mixed-Type Data

Alexander Marx and Jilles Vreeken

Max Planck Institute for Informatics and Saarland University, Saarbrücken, Germany  
`{amarx,jilles}@mpi-inf.mpg.de`


## A Appendix

In the appendix we give additional results and describe the multivariate cause-effect pairs and their corresponding datasets in more detail.

### A.1 Encoding of the internal nodes

### A.2 Synthetic data

Here we show the additional plot about how both scores deal with dimensionality, i.e. when the number of dimensions  $k = l$  increases. In Figure 1 we show the results for  $k, l \in \{2, 3, 5, 7, 11\}$  on nominal, numeric and mixed-type data for  $\text{CRACK}_\Delta$  and  $\text{CRACK}_\delta$ .  $\text{CRACK}_\delta$  performs better on mixed-type data and is equally good on the single-type data sets. In general, both approaches work well in high dimensions.



**Fig. 1.** Accuracy of  $\Delta$  (left) and  $\delta$  (right) on symmetric dimensions  $k \in \{2, 3, 5, 7, 11\}$  for nominal, numeric and mixed-type data.

**Data sets** *Haberman* is a data set on medical case studies describing the survival of patients who had undergone surgery for breast cancer between 1958 and 1970 [3].  $X$  consists of the age of the patient at time of operation, the patient’s year of operation and the number of positive axillary nodes detected.  $Y$  is the survival status, which is binary and divided into longer or at most five years ( $X \rightarrow Y$ ). The *Iris* data set contains data about three types of the Iris plant

( $Y$ ) and four features dependent on which the type can be determined [1]. Next, we extract four cause-effect data sets from the *Mammals* data set [4], which consists of both climate data and presence records of 121 mammal species over 2183 areas of  $50 \times 50$ km in Europe. We assume that elevation, precipitation, average temperature and the annual temperature range ( $X$ ) cause the presence of a mammal and not contrarily. We created three data sets, *Canis*, *Lepus* and *Martes*, each containing locations of different types of the named species and one data set containing all three of them. Last, we created a data set based on the octet data set [2,6]. Marx and Vreeken [5] created 10 univariate cause effect pairs based on the data set that had all the same effect that we combined to a single multivariate data set.

## References

1. R. A. Fisher. The use of multiple measurements in taxonomic problems. *Ann Eugen*, 7(2):179–188, 1936.
2. L. M. Ghiringhelli, J. Vybiral, S. V. Levchenko, C. Draxl, and M. Scheffler. Big data of materials science: Critical role of the descriptor. *PRL*, 114, 2015.
3. S. J. Haberman. Generalized residuals for log-linear models. In *Proceedings of the 9th international biometrics conference*, pages 104–122, 1976.
4. H. Heikinheimo, M. Fortelius, J. Eronen, and H. Mannila. Biogeography of European land mammals shows environmentally distinct and spatially coherent clusters. *J. Biogeogr.*, 34:1053–1064, 2007.
5. A. Marx and J. Vreeken. Telling Cause from Effect by MDL-based Local and Global Regression. In *ICDM*, pages 307–316. IEEE, 2017.
6. J. A. Van Vechten. Quantum dielectric theory of electronegativity in covalent systems. i. electronic dielectric constant. *Physical Review*, 182(3):891, 1969.